



2012 River Herring Stock Assessment

Dr. John Sweka, *Chair*, U.S. Fish and Wildlife Service

Dr. Michael Bailey, U.S. Fish and Wildlife Service

Michael Brown, Maine Department of Marine Resources

Kiersten Curti, National Marine Fisheries Service

Dr. Katie Drew, Atlantic States Marine Fisheries Service

Phil Edwards, Rhode Island Department of Environmental Management

Kathryn Hattala, New York Department of Environmental Conservation



Andrew Kahnle, New York Department of Environmental Conservation

Laura Lee, North Carolina Division of Marine Fisheries

Dr. Gary Nelson, Massachusetts Division of Marine Fisheries


Robert Sadzinski, Maryland Department of Natural Resources

Kate Taylor, Atlantic States Marine Fisheries Commission






Outline



- Overview of state & regional data sets
- Coast-wide comparisons & trend analysis
 - Total landings & incidental catch
 - Biological data
 - Total mortality estimates
- Stock assessment models
- Conclusions





Data Overview

- **57 “systems” on Atlantic coast**
 - 9 FI & FD data categories
- **54% are blank**
- **Only 26% have complete or “good” data**
 - Most occurs in NE states

State	River	Time series	By species	Harvest	Age	Length	Weight	Repeat Spawner	FI Adult	FI JAI	FD CPUE
ME	Damariscotta	1943-2010		•							
	St. George	1943-2010		•							
	Union	1975-2010		•							
	Orland	1943-2010		•							
	Androscooggin	1983-2010	•		•	•					
	Sebasticook	2000-2010		•							
	Merrymeeting Bay/Tribs	1979-2009	•							•	
	Gulf of Maine	2000-2010	•						•		
NH	Exeter/Squamscott	1991-2010	•	•	•	•		○	•		
	Lamprey	1991-2010	•	•	•	•		○	•		
	Winnicut	1991-2010	•	•	•	•		○	•		
	Oyster	1991-2010	•	•	•	•		○	•		
	Cocheco	1991-2010	•	•	•	•		○	•		
	Taylor	1991-2010	•	•	•	•		○	•		
	Great Bay Estuary	1997-2010	x			x					x
MA	Mattapoissett	1988-2010	•	•	○	○	○		•		
	Monument	1980-2010	•	•	○	○	○	○	•		
	Nemasket	1996-2010	•	•	○	○	○				
	Parker	1971-1978, 2000-2010	•	•	○	○			•		
	Town	2000-2010		•					•		
	Agawam	2006-2010			○	○					
	Back	2007-2010	•	•					•		
	Charles	2008-2009		•	•	•	•	•	•		
	Mystic	2004-2010	•	•			•	•			
	Quashnet	2004	•					•			
	Stony Brook	1978-2004	•		○	○	○	○	○		
RI	Gilbert Stuart	1981-2010		•	•	•	•	•	•	○	
	Nonquit	1999-2010		•	•	•	•	•	•	○	
	Buckeye Brook	2003-2010							•		
	Pawcatuck	1988-2010			x	x	x	x	○	•	
	Ocean waters	1979-2010				•			•	•	
	Naragansett Bay	1988-2010				•			•	•	
	Coastal ponds	1992-2010							•	•	
CT	Bride Brook	1966-1967, 2003-2011	•			○			•		
	Connecticut River	1975-2011	•			○				○	
	Farmington River	1976-2011	•						•		
	Thames River	1996-2011	•						•		
NY	Hudson	1975-2010	•	○	○	○		○	○	○	○
DE, NJ, PA	Delaware River	1980-2010	○	○		○			○	○	○
	Delaware Bay	1966-2010	○	○						○	○
MD	Nanticoke	1959-2010	○		○	○		○		○	○
	Susquehanna	1972-2010	○						x		
	Chesapeake Bay	1959-2010			○						○
MD, VA, DC	Potomac River	1959-2010		•		○			○	○	○
VA	James	1966-2010	○	•	○	○	○	○	○	○	○
	Rappahannock	1966-2010	○	•	○	○	○	○	○	○	○
	York	1966-2010	○	•	○	○	○	○	○	○	○
NC	Albemarle Sound	1972-2009		○				○	○	•	
	Chowan River	1972-2009	•	•	•	•	•	○			•
SC	Wynah Bay									x	
	Santee-Cooper	1969-2010	○	•	○	○	○	○	○	x	•
	Savannah River									x	
	Ashley-Combahee-Edisto Basin									x	
GA	Altamaha River	2010								x	
	Ogeechee River	2010								x	
	Savannah River	2010								x	
FL	St. John's River	2001-2010	•			•			•	○	





Reported US Landings






Reported NAFO Landings






Total annual incidental catch by species







Biological Data



- Observers also collect data on length frequency of incidental catch
- Incidental catch included small fish of size classes not observed in river samples






Trend Analysis

- Commercial CPUE (NY, NJ, MD, PRFC, VA, NC, SC)
- Run Sizes (ME, NH, MA, RI, CT)
- YOY Indices (ME, RI, CT, NY, NJ, MD, DC, VA, NC)
- Trawl Surveys (NEFSC, MA, RI, CT, NJ, NC)
- Mean length and length-at-age (ME, NH, MA, RI, NY, MD, NC, SC, FL)
- Repeat Spawners (ME, NH, MA, RI, NY, MD, NC, SC)
- Total Mortality (ME, NH, MA, RI, MD, NC, SC)



Commercial CPUE



Alewife:


3 of 4 series showed historical declines and some increases in recent years.

Blueback:

2 of 3 series have declined or are showing declines in recent years.

Combined Species:

3 of 4 series have declined.




Run Sizes

- Run sizes for alewife, blueback herring and combined species showed historical and recent (1999-2007) declines in abundance.
- Alewife/Combined Species run sizes in eight of nine NE rivers (with long time series: 1984-2010) showed historical declines (mid 1990s or after 1999-2000) but have increased in the last 2-3 years.
- Blueback run sizes in two of two NE rivers declined over time (as early as 1985).
- Population sizes in Chowan River, NC declined precipitously after 1985 and abundance remains low.



YOY FI Surveys



➤ For recent years (2000-2007):

- Alewife:

- ➔ **3 indices declining**

- ➔ 3 indices showed no trend

- ➔ **1 index increasing**

- Blueback from eight rivers showed:


- ➔ 4 rivers showed no trend

- ➔ **4 rivers declining**

➤ Similar patterns among indices from the same region



FI Trawl Surveys



➤ Alewife


- **Increasing trend: 4 surveys**
- **Flat/no trend: 3 surveys**
- **Decreasing trend: 1 survey**

➤ Blueback herring

- **Increasing trend: 2 surveys**
- **Flat/no trend: 4 surveys**
- **Decreasing trend: 2 survey**




Mean Length



➤ Mean length of male and female alewife and blueback herring have **declined** over time by 13-45 mm TL in 7 of 13 rivers examined

➤ **Significant declines for rivers with long time series**






Maximum Age Observed



- Alewife maximum age has **declined** by 1 to 2 ages in MA, RI, MD and NC. Trends in ME and NH have been stable or **increasing**.
- Blueback maximum age has **declined** by 1 to 4 ages in MA, MD and NC. Trends in NH have been stable or **increasing**.




Length-at-Age





➤ Significant **declines** in mean length for one or more ages in:

- Alewife – ME, NH, RI, MD and NC
- Blueback – NH, MD, NC






Repeat Spawner Data



- Data available from 12 rivers
- Few datasets with time series
- **Declining** trends in the **Gilbert-Stuart River (RI)** for combined species and **Nanticoke River** for blueback herring only
- **No or little trends in the remaining rivers**



Total Mortality (Z) Estimates




- **Developed from observed age-structure**
- **Chapman-Robson – least biased estimator**
- **At least 3 age classes must be present**




Z Benchmarks

- **Total mortality was high for all stocks examined**
- **Three year average of observed Z values were **above** the $Z_{20\%SPR}$ benchmark for 12 of the 18 stocks.**
- **Three year average of observed Z values were **between** the $Z_{40\%SPR}$ and $Z_{20\%SPR}$ benchmarks for the remaining 6 stocks.**



Coastwide Model

- Depletion-Based Stock Reduction Analysis (**DB-SRA**)
- Developed on the west coast to generate management parameters (e.g., MSY) for data-poor species
- SASC & Peer Review Panel had concerns about model structure and assumptions and recommends further development




Conclusions

➤ **River herring have declined coast-wide**

- Declining commercial landings following the 1960s
- Declining commercial CPUE
- Declining run counts in many rivers
- Declines in average length and size-at-age in many rivers
- SCAA and DB-SRA model runs


➤ **Fisheries independent indices were quite variable**

- Most started after the decline in commercial landings
- Currently observing relatively small amounts of inter-annual variation
- Regional (north vs. south) patterns may be due to climate change



Stock Status

- **The coastwide meta-complex of river herring on the US Atlantic coast is depleted to near historic lows**
- “Depleted” status indicates that there was evidence for declines in abundance due to a number of factors, but the relative importance of these factors in reducing river herring stocks could not be determined.



Stock Status

➤ **52 in-river stocks** for which data were available

- Historically:

- **22 were depleted**

- **1 stock was increasing**

- **28 stocks could not be determined**


- In most recent years:

- **2 were increasing**

- **4 were decreasing**

- **9 were stable**

- **38 rivers did not having enough data**



Stock Status

- **Overfished and overfishing status could not be determined for the coastwide stock complex**
- **Management actions to reduce total mortality are needed.**
- **Recovery of river herring stocks will need to address multiple factors (e.g., fish passage, predation, water quality, climate change, etc.) in addition to harvest.**